


This article was downloaded by: [University College London]

On: 10 February 2014, At: 09:41

Publisher: Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Social Neuroscience

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/psns20>

Distinct effects of self-construal priming on empathic neural responses in Chinese and Westerners

Chao Jiang^a, Michael E. W. Varnum^b, Youyang Hou^a & Shihui Han^a

^a Department of Psychology, PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, China

^b Department of Psychology, Arizona State University, Phoenix, AZ, USA

Published online: 16 Dec 2013.

To cite this article: Chao Jiang, Michael E. W. Varnum, Youyang Hou & Shihui Han (2014) Distinct effects of self-construal priming on empathic neural responses in Chinese and Westerners, *Social Neuroscience*, 9:2, 130-138, DOI: [10.1080/17470919.2013.867899](https://doi.org/10.1080/17470919.2013.867899)

To link to this article: <http://dx.doi.org/10.1080/17470919.2013.867899>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Distinct effects of self-construal priming on empathic neural responses in Chinese and Westerners

Chao Jiang¹, Michael E. W. Varnum², Youyang Hou¹, and Shihui Han¹

¹Department of Psychology, PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, China

²Department of Psychology, Arizona State University, Phoenix, AZ, USA

The present study investigated whether and how self-construal priming influences empathic neural responses to others' emotional states. We recorded event-related brain potentials to stimuli depicting the hands of unknown others experiencing painful or non-painful events from Chinese and Western participants after they had been primed in three conditions (independent self-construal priming, interdependent self-construal priming, and a control condition). Stimuli depicting painful events (as opposed to non-painful ones) elicited a positive shift of the fronto-central activity at 232–332 ms and of the central-parietal activity at 440–740 ms in the control condition. Moreover, neural responses to stimuli depicting painful (vs. non-painful) situations at 232–332 ms were decreased by interdependent self-construal priming among Chinese and by independent self-construal priming among Westerners. Our findings suggest that self-construal priming modulates sensitivity to perceived pain in unknown others and that this effect varies with culture.

Keywords: Culture; Empathy; ERP; Self-construal; Pain.

Self-construal refers to how individuals define and make meaning of the self and how one does so has a number of psychological and behavioral consequences (Cross, Hardin, & Gercek-Swing, 2011). It has been well documented that self-construals vary across cultures such that the self is viewed as bounded and autonomous (an independent self-construal) in Western cultures but is seen as interconnected and overlapping with close others (an interdependent self-construal) in East Asian cultures (Markus & Kitayama, 1991). Different self-construals tend to covary with a suite of cognitive and affective tendencies (Varnum, Grossmann, Kitayama, & Nisbett, 2010), and making different self-construals salient leads to changes in these psychological processes (Oyserman & Lee, 2008) and in neural responses (Han & Northoff, 2008; Han et al., 2013).

For example, neuroimaging studies have shown evidence that self-construal priming (a procedure that asks participants to read essays) with independent (e.g., “I”, “mine”) or interdependent pronouns (e.g., “we”, “ours”), in order to shift their self-construals toward independence or interdependence (Gardner, Gabriel, & Lee, 1999; Oyserman & Lee, 2008), affects neural substrates underlying self-related processes. Wang et al. recorded event-related potentials (ERPs) to painful and non-painful electrical stimulations from adults after self-construal priming (Wang, Ma, & Han, in press). They found that independent (as opposed to interdependent) self-construal priming increased an early somatosensory activity over the frontal/central region in response to painful stimulation applied to participants' hands. Neural responses to images of one's own face are also modulated by

Correspondence should be addressed to: Shihui Han, Department of Psychology, Peking University, 5 Yiheyuan Road, Beijing 100871, People's Republic of China. E-mail: shan@pku.edu.cn

This work was supported by the National Natural Science Foundation of China [Project 30910103901, 91024032, 81161120539; 91224008 and the National Basic Research Program of China [973 Program 2010CB833903].

self-construal priming such that independent (as opposed to interdependent) self-construal priming increases the right frontal activity during perception of one's own face (Sui & Han, 2007).

There is also evidence suggesting that self-construal priming affects neurocognitive processes related to others. For example, a recent transcranial magnetic stimulation (TMS) study found that interdependent self-construal primes superimposed throughout videos depicting others' hand movements (e.g., squeezing a rubber ball) increased motor-evoked potentials recorded from a participant's hand relative to a no-priming baseline condition (Obhi, Hogeveen, & Pascual-Leone, 2011), suggesting self-construal-induced changes in the motor system that may reflect self-construal effects on behavioral mimicry in social settings. In another study, Harada et al. found that the dorsal medial prefrontal cortex showed increased activity during implicit evaluation of information related to one's father (vs. a stranger) when bicultural (i.e., Asian-American) individuals were primed with independent (but not interdependent) self-construals (Harada, Li, & Chiao, 2010). Further, a recent study with Chinese young adults found that priming an interdependent self-construal increased activity in the bilateral ventral striatum in response to winning money for a friend (Varnum, Shi, Chen, Qiu, & Han, in press). Taken together, these findings suggest that self-construal priming modulates neural activity involved in cognitive and affective processing of information about the self and close others.

The present study examined whether self-construal priming modulates brain activity underlying the processing of strangers' emotional states. Specifically, we investigated the effect of self-construal priming on the neural activity elicited by perceived pain in others. We recorded ERPs from healthy adults when they perceived painful versus non-painful stimuli applied to hands of unknown others. Recent ERP studies have shown that perception of painful versus non-painful stimuli applied to others' body parts (e.g., hands) induces positive shift of ERP amplitudes in a large time window (from 140 to 660 ms after stimulus onset) over the frontal/central/parietal regions (Decety, Yang, & Cheng, 2010; Fan & Han, 2008; Han, Fan, & Mao, 2008; Li & Han, 2010). Viewing others' pain versus neutral expressions also elicits increased positivity at 128–188 ms poststimulus over the frontal regions (Sheng & Han, 2012; Sheng, Liu, Zhou, Zhou, & Han, 2013). In addition, the amplitudes of these neural responses are correlated with subjective reports of the degree of others' pain and with subjective reports of one's own discomfort (Fan & Han, 2008; Li & Han, 2010; Sheng & Han, 2012).

Thus, these neural responses are associated with understanding and sharing others' pain (or empathy for others' pain). We were interested in whether self-construal priming that temporarily highlights different self-construals (e.g., independence or interdependence) modulates differential ERP amplitudes elicited by painful versus non-painful stimuli.

In the current study, we recruited both Chinese and Westerners for two reasons. First, it is well documented that interdependent self-construals are predominant in China (de Greck et al., 2012; Li, Zhang, Bhatt, & Yum, 2006; Ma et al., in press), whereas independent self-construals are predominant in Western cultures (Thomsen, Sidanius, & Fiske, 2007, see Markus & Kitayama, 1991; Oyserman, Coon, & Kimmelmeier, 2002 for review). Second, in a recent ERP study we found that priming interdependent self-construals decreased a frontal/central activity to one's own face for British participants, whereas priming independent self-construals suppressed this activity to a friend's face for Chinese participants (Sui, Hong, Liu, Humphreys, & Han, 2013). This finding suggests that the effects of self-construal priming on neural responses may be constrained by participants' long-term cultural experiences. Thus, we were interested in whether self-construal priming produces similar effects on empathic neural responses in individuals from different cultures. As the previous research has shown that empathic neural responses in male and female participants are differentially sensitive to attitude toward others and task modulations (Han et al., 2008; Singer et al., 2006), the current work only tested females in order to exclude potential effects of gender differences in empathic neural responses.

METHODS

Participants

Eighteen Chinese females (aged between 19 and 26 years, mean = 20.50, SD = 1.46, all native Chinese speakers and right-handed) and eighteen Westerner females (aged between 19 and 31 years, mean = 21.83, SD = 3.17, 16 native English speakers, 1 native German speaker, 1 native Spanish speaker, 16 right-handed) participated in the study as paid volunteers. All participants were undergraduate or graduate students at the time of testing. Chinese participants had not spent any time outside China and Western participants had been in China for less than 1 month. Chinese participants were proficient in English and Western participants had not studied Chinese at the time of testing. All

participants had normal or corrected-to-normal vision and were not color-blind. Informed consent was obtained prior to the study. This study was approved by a local ethic committee at the Department of Psychology, Peking University.

Self-construal priming

The materials for self-construal priming consisted of six short essays that described trips to the countryside and were used in our prior studies (Lin & Han, 2009; Sui & Han, 2007; Sui et al., 2013). Each essay consisted of 300–350 words and was presented in Chinese for Chinese participants and in English for Western participants. The priming materials contained independent pronouns (e.g., “I”, “mine”) during independent self-construal priming and interdependent pronouns (e.g., “we”, “ours”) during interdependent self-construal priming. The materials used for the control priming did not contain either type of pronoun. After reading each essay, participants had to indicate the number of occurrences of independent (or interdependent) pronouns or specific nouns (e.g., “lake,” “park”).

Visual stimuli

Visual stimuli consisted of 40 color pictures showing hands in painful situations and 40 color pictures of hands in non-painful situations (illustrated in Figure 1), similar to those used in our previous work (Fan & Han, 2008; Gu & Han, 2007). The pictures were shot from the first-person perspective and described accidents that may happen in everyday life. Painful stimuli included situations such as a hand trapped in a door or cut by scissors. The stimuli were presented in the center of a grey background on

a 21-inch color monitor. Each stimulus subtended a visual angle of $2.58^\circ \times 3.43^\circ$ (width \times height) at a viewing distance of 100 cm. A pilot test of Chinese and Caucasian participants suggested that they were unable to tell whether hands in pictures were from Asian or Caucasians.

Procedure

Participants were presented with images of others' hands receiving painful and non-painful stimuli while an electroencephalogram (EEG) was recorded. There were 9 blocks of 80 trials for each participant. After reading one essay, EEG was recorded during three blocks of trials. The order of reading essays for independent, interdependent, and control priming was counterbalanced across participants. Each block started with the presentation of instructions for 3 s followed by 80 trials. On each trial, a painful or non-painful stimulus was presented for 200 ms at the center of the screen, which was followed by a fixation cross with a duration varying randomly between 800 and 1600 ms. Painful and non-painful stimuli were presented in a random order. Participants were asked to judge whether or not the target felt pain by a button press using the left or right index finger. The assignment of the left or right index finger to the painful and non-painful stimuli and the order of the priming procedure were counterbalanced across participants. Participants completed the Interpersonal Reactivity Index (IRI) to measure their trait-level empathy (Davis, 1983) after EEG recording. ERP recording and analysis were similar to those in our previous work (Fan & Han, 2008). Mean amplitudes of each ERP component were calculated at frontal (Fz, FCz, F3–F4, FC3–FC4), central (Cz, CPz, C3–C4, CP3–CP4), and parietal (Pz, P3–P4) electrodes.



Figure 1. Illustration of a painful and a non-painful stimulus used in the current study.

RESULTS

Behavioral performance

Both behavioral and ERP data were subject to a $3 \times 2 \times 2$ repeated-measures analyses of variance (ANOVA) with Priming (independent, interdependent, or control) \times Pain (Painful vs. Non-painful) as within-subjects variables and Group (Westerners vs. Chinese) as a between-subjects variable. The ANOVA of reaction times (RTs) only showed a significant main effect of Pain, as participants responded faster to painful than to non-painful stimuli, $F(1, 34) = 10.343$, $p < .005$. Analysis of response accuracy data did not reveal any significant effects, $ps > .1$ (Table 1). There were no group differences on IRI subscales with the exception of personal distress. Chinese reported greater personal distress ratings compared to Westerners, $t(1,34) = -3.651$, $p < .001$ (Table 2).

ERP results

Grand-averaged ERPs to painful and non-painful stimuli are illustrated in Figure 2. ANOVAs of the mean ERP amplitudes at 232–332 ms, which covered both the N2 and N320 components, showed a significant main effect of Pain at fronto-central electrodes, ($F_s(1,34) = 7.28\text{--}24.36$, $ps < .01$), such that painful stimuli induced a positive shift of ERP amplitude relative to non-painful stimuli. Importantly, there were significant three-way—Group \times Priming \times Pain—

interactions observed at fronto-central electrodes ($F_s(2,68) = 3.13\text{--}5.55$, $ps < .05$), indicating that priming independence decreased empathic responses for Westerners (relative to other conditions), whereas priming interdependence decreased empathic responses (relative to other conditions) for Chinese participants.

Next we analyzed the effects of priming separately for each cultural group. For Chinese participants, the ANOVAs of the ERP amplitude at 232–332 ms showed a significant main effect of Pain ($F_s(1,17) = 4.748\text{--}9.049$, $ps < .05$) and a significant interaction of Pain \times Priming at fronto-central electrodes ($F_s(2, 34) = 3.67\text{--}3.74$, $ps < .05$). Post-hoc t -tests confirmed that, relative to non-painful stimuli, painful stimuli elicited a significant positive shift of the ERP amplitudes in this time window in the independent self-construal priming ($t_s(17) = 2.50\text{--}3.10$, $ps < .05$) and control priming ($t_s(17) = 2.30\text{--}2.47$, $ps < .05$) conditions but not in the interdependent self-construal priming condition ($ps > .1$).

For Westerners, ANOVAs of ERP amplitudes at 232–332 ms revealed a significant main effect of Pain ($F_s(1,17) = 14.765\text{--}16.184$, $ps < .01$) and a significant interaction of Pain \times Priming at frontal and parietal electrodes ($F_s(2, 34) = 3.46\text{--}4.16$, $ps < .05$). Post-hoc t -tests confirmed that painful compared to non-painful stimuli significantly elicited positive shifts of the ERP amplitudes in this time window in the interdependent self-construal priming ($t(17) = 3.40\text{--}4.04$, $ps < .005$) and control priming ($t(17) = 3.21\text{--}3.96$, $ps < .005$) conditions but not in the independent self-construal priming condition ($ps > .5$).

TABLE 1
Mean RTs and response accuracy (standard deviation) in each condition

	RTs (ms)			Accuracy (%)		
	Control	Independent	Interdependent	Control	Independent	Interdependent
<i>Westerners</i>						
Painful	600(80)	588(91)	616(90)	79.6(10)	79.8(10.4)	79.0(8.3)
Non-painful	619(79)	616(97)	645(96)	86.4(14)	83.1(16.9)	83.6(11)
<i>Chinese</i>						
Painful	628(71)	649(98)	628(63)	83.7(8.9)	79.1(16.7)	82.1(8.2)
Non-painful	641(73)	671(87)	646(63)	86(11)	79.5(13.7)	85.4(8.0)

TABLE 2
Scores on IRI subscales of each group

	Westerners	Chinese
Perspective-taking scale	20.4(4.46)	21.0(2.93)
Fantasy scale	15.0(5.32)	17.4(2.96)
Empathic concern scale	18.6(5.56)	20.6(2.61)
Personal distress scale	11.4(6.03)	17.6(3.99)

Note: Personal distress ratings were significantly higher in Chinese than Westerners.




Figure 2. Illustration of ERP results in the current study. ERPs to painful and non-painful stimuli recorded at electrode F4 are illustrated in (a) the control priming condition, (b) independent self-construal priming condition, and (c) interdependent self-construal priming condition. The grey bars indicate the time window in which the ANOVA of mean ERP amplitudes showed significant 3-way interactions. (d) The mean amplitudes and standard deviations of difference waves to painful vs. non-painful stimuli at 232–332 ms post-stimulus at electrode F4 are illustrated in different priming conditions. (e) The voltage topographies illustrate the scalp distributions of the difference wave to painful vs. non-painful stimuli at 232–332 ms poststimulus in each priming condition. ** $p < .01$.

There was a main effect of Pain on mean P3 amplitudes at 440–740 ms ($F_s(1,34) = 20.87\text{--}37.11$, $ps < .001$). However, there was no significant interaction between Group, Priming, and Pain ($ps > .1$). Thus, self-construal priming did not significantly affect empathic neural responses in this time window.

Finally, to examine whether an individual's trait empathy can predict the effect of self-construal priming on empathic neural responses, we calculated the difference in neural responses at 232–332 ms to

images of others' hands receiving painful versus non-painful stimuli between independent (or interdependent) self-construal priming and the control conditions across all participants. We then examined the correlation between IRI scores and the differential ERP amplitudes in this time window. However, correlations were not significant ($ps > .5$), suggesting that the effect of self-construal priming on empathic neural responses does not vary significantly as a function of individuals' trait empathy.

DISCUSSION

The current work examined the effect of self-construal priming on empathic neural responses to others' pain. Our ERP results in the control condition showed that painful stimuli elicited a positive shift in neural activity over the fronto-central regions in an early time window and over the parietal region in a later time window in both Chinese and Westerners. These results replicate previous findings (Decety et al., 2010; Fan & Han, 2008; Han et al., 2008; Li & Han, 2010) and suggest that, for both cultural groups, there are similar default neural responses to perceived pain in unknown individuals. Moreover, we found that temporarily highlighting different self-construals (e.g., interdependence or independence) significantly influenced neural responses to others' suffering. Social cognition consists of two components, i.e., processing information about the self and processing information about others (Iacoboni, 2006; Sedikides & Skowronski, 2009). Our ERP results showed that the manner in which one conceptualizes the self influences one's neural responses to others' pain. The effect of self-construal priming was limited to fronto-central activity at 232–332 ms, which is associated with the automatic component of empathy (Fan & Han, 2008). Thus, our ERP results suggest that self-construal priming modulates the early automatic component of empathy but produces little effect on the later controlled component of empathy in the P3 time window (Fan & Han, 2008).

Interestingly, we found that the effects of independent and interdependent self-construal priming on empathic neural responses significantly differed between Chinese and Westerners. Relative to the control condition, independent self-construal priming decreased empathic neural responses among Westerners, whereas interdependent self-construal priming decreased empathic neural responses among Chinese. Although chronic and temporary self-construals have been found to have parallel effects in terms of variables like cognitive style (Oyserman & Lee, 2008; Sui, Liu, & Han, 2009; Sui et al., 2013; Varnum et al., 2010), our findings hint that this may not be the case for empathic neural responses. Our findings also suggest that social contexts that promote chronic views of the self as independent or interdependent do not have a direct effect on empathic neural responses to strangers but may interact with temporarily activated self-views to modulate empathic neural responses.

One possible reason for the distinct effects of self-construal priming observed in Chinese and

Westerners is the default ways in which the self is construed in Chinese and Western societies. One predominant theory holds that the independent self-construal dominates Western cultures and the interdependent self-construal dominates East Asian cultures (Markus & Kitayama, 1991). Moreover, interdependent self-construals induce a strong boundary between in-group (including the self and close others) and out-group (non-close others such as strangers), whereas independent self-construals define a strong boundary between the self and any others (including close and non-close others) (Markus & Kitayama, 2010). In cultures where the self is constituted of no one but the individual (such as the US or Western Europe), activating an independent mind-set may cause all others to be excluded from the self and thus reduce empathic responses to out-group members (i.e., unknown others). In cultures where the self is constituted of the individual and close others such as family members and friends (such as China), activating an interdependent mind-set may enhance the boundary between in-group (i.e., self and close others) and out-group (i.e., unknown others), which in turn may weaken empathic neural responses to perceived pain in unknown others. This account is in line with a recent finding that Asian-Americans (who tend to have a more interdependent view of the self) showed less empathic accuracy when determining strangers' emotions but greater accuracy when determining friends' emotions compared with European Americans (who tend to have a more independent view of the self; Ma-Kellams & Blascovich, 2012). This account is also consistent with recent findings that neural responses to out-group members' pain are significantly reduced compared to responses to in-group members' pain (Avenanti, Sirigu, & Aglioti, 2010; Hein, Silani, Preuschoff, Batson, & Singer, 2010; Sheng & Han, 2012; Sheng et al., 2013; Xu, Zuo, Wang, & Han, 2009; Sheng, Liu, Li, Fang, & Han, in press).

Previous studies have shown that empathic neural responses are modulated by a host of factors such as top-down attention to painful cues (Gu & Han, 2007), personal experiences (Cheng et al., 2007), affective connection (Singer et al., 2006), morality salience (Luo et al., in press), socioeconomic status (Ma, Wang, & Han, 2011), and intergroup experiences (Xu et al., 2009; Zuo & Han, 2013). The fact that the two cultural groups in our study did not show differences in empathic neural responses in the control condition suggests that chronic self-construals may not be one of these factors. However, as the present study did not measure the chronic self-construal of

participants, the relationship between chronic self-construal and empathy remains a matter for further investigation.

One possible alternative interpretation of our results is that different routes to reduced empathy are more easily activated in different cultural contexts. Previous research suggests that while some degree of self–other overlap is necessary for empathy to occur, a strong sense of self–other overlap may actually decrease empathy (Preston & Hofelich, 2012), as may extreme self-focus (Biscardi & Schill, 1985; Konrath, Bushman, & Grove, 2009; Watson, Grisham, Trotter, & Biderman, 1984). Thus, it is possible that priming interdependence may activate an overly strong sense of self–other overlap in Chinese participants, whereas priming independence activated an overly strong sense of self-focus among Westerners. As we did not assess participants' subjective feeling of overlap with the targets, it is unclear whether the effects of interdependence priming on Chinese participants' empathic responses were due to either a greater sense of excluding the target individuals from the self or due to a greater sense of including them in the self. Although we suspect the former interpretation is more likely as interdependent self-construal involves a less rather than more permeable boundary between the self and non-close others (Falk, Heine, Yuki, & Takemura, 2009; Markus & Kitayama, 2010), future research should test these possible accounts by directly measuring participants' chronic self-construals.

Neither priming interdependent self-construals in Westerners nor priming independent self-construals in Chinese produced significant influences on empathic neural responses to others' suffering relative to the control priming. Recently, Sui et al. (2013) primed British and Chinese participants with independent and interdependent self-construals during perception of one's own face or a friend's face. They found that priming an interdependent self-construal reduced an anterior N2 to one's own face for British, whereas priming an independent self-construal suppressed the anterior N2 to a friend's face for Chinese. These results together suggest that priming both culturally congruent and incongruent self-construals can affect neural substrates involved in social cognition. However, the pattern of effects depends on stimuli and task demands. To our knowledge, the previous studies have shown evidence that self-construal priming modulates frontal activity related to self-face recognition (Sui & Han, 2007), occipital activity related to visual perception (Lin, Lin & Han, 2009), motor activity (Obhi et al., 2011), and somatosensory activity related to physical pain (Wang et al., in press).

These findings suggest that priming interdependence may facilitate neural activity related to the processing of others, whereas priming independence may enhance neural activity related to the processing of oneself. However, the results of our previous (Sui et al., 2013) and current studies suggest that self-construal priming may not generate a culturally universal effect on human brain activity. Instead, one's self-construals formed during long-term cultural experiences may constrain the effect of self-construal priming.

The previous ERP study found that the P2 amplitude at 128–188 ms in responses to pain versus neutral expression was positively correlated with rating scores of empathic concern (Sheng & Han, 2012), suggesting an association between trait empathy and empathic neural activity in this time window. In the current work, we found that the effect of self-construal priming on empathic neural responses in the N2/N320 time window was not correlated with individuals' trait empathy. Empathic neural responses in the control condition did not differ between the two cultural groups even though self-report personal distress was higher in Chinese compared to Westerners. Thus, the neural responses in this time window may not be associated with an individual's trait empathy.

We should note some important limitations of the current work. First, although participants were from societies that tend to differ in the type of self-construal which is predominant, we did not measure participants' self-construals. Therefore, although our ERP results revealed distinct effects of independent and interdependent self-construal priming on the empathic neural responses of Chinese and Westerners, it is still unclear whether these effects were linked to group differences in chronic self-construal. Thus future research should further examine how self-construal priming interacts with one's chronic self-construal to modulate empathy for others' pain. This would allow future researchers to determine whether the interaction we observed was in fact driven by chronic cultural group differences in self-construal. Second, previous fMRI studies have shown that perceiving painful stimulation applied to others' body parts activated several brain regions including the anterior cingulate cortex (ACC) and the insula (e.g., Gu & Han, 2007; Jackson, Meltzoff, & Decety, 2005). The current ERP results did not identify the source of the increased fronto-central activity at 232–332 ms in response to others' pain. Thus, it remains unclear whether the effect of self-construal priming on empathic neural responses originated in the ACC, the insula, or other brain regions. Finally, as the current work only

recorded ERPs in response to others' hands, it remains unknown whether self-construal priming modulates neural responses to others' facial expressions of pain in a way similar to what we observed here. These questions can be clarified in future research.

In sum, our findings shed new light on the relationship between self-construals and empathy for pain. Together with previous research (e.g., Sui et al., 2013), our findings suggest that self-construal priming may produce significant effects on neurocognitive processes of the self and others, and these effects may vary across cultures (e.g., Chinese and Westerners). Future research may investigate what effects such priming might have on empathy for close others, given that the previous study of Chinese showed differential neural activity in response to witnessing a friend or a stranger experiencing social pain (Meyer et al., 2013). It would also be interesting to examine the effect of self-construal priming on empathy for other emotions since de Greck et al. (2012) have found that culture influences neural correlates of empathy for anger.

Original manuscript received 3 March 2013

Revised manuscript accepted 17 November 2013

First published online 16 December 2013

REFERENCES

- Avenanti, A., Sirigu, A., & Aglioti, S. M. (2010). Racial bias reduces empathic sensorimotor resonance with other-race pain. *Current Biology*, *20*, 1018–1022.
- Biscardi, D., & Schill, T. (1985). Correlations of narcissistic traits with defensive style, machiavellianism, and empathy. *Psychological Reports*, *57*, 354.
- Cheng, Y., Lin, C. P., Liu, H. L., Hsu, Y. Y., Lim, K. E., Hung, D., & Decety, J. (2007). Expertise modulates the perception of pain in others. *Current Biology*, *17*, 1708–1713.
- Cross, S. E., Hardin, E. E., & Gercek-Swing, B. (2011). The what, how, why, and where of self-construal. *Personality and Social Psychology Review*, *15*, 142–179.
- Davis, M. H. (1983). The effects of dispositional empathy on emotional reactions and helping: A multidimensional approach. *Journal of Personality*, *51*, 167–184.
- Decety, J., Yang, C., & Cheng, Y. (2010). Physicians down-regulate their pain empathy response: An event-related brain potential study. *NeuroImage*, *50*, 1676–1682.
- de Greck, M., Shi, Z., Wang, G., Zuo, X., Yang, X., Wang, X., . . . , Han, S. (2012). Culture modulates brain activity during empathy with anger. *NeuroImage*, *59*, 2871–2882.
- Falk, C. F., Heine, S. J., Yuki, M., & Takemura, K. (2009). Why do Westerners self-enhance more than East Asians? *European Journal of Personality*, *23*, 183–203.
- Fan, Y., & Han, S. (2008). Temporal dynamic of neural mechanisms involved in empathy for pain: An event-related brain potential study. *Neuropsychologia*, *46*, 160–173.
- Gardner, W. L., Gabriel, S., & Lee, A. Y. (1999). 'I' value freedom, but "we" value relationships: Self-construal priming mirrors cultural differences in judgment. *Psychological Science*, *10*, 321–326.
- Gu, X., & Han, S. (2007). Attention and reality constraints on the neural processes of empathy for pain. *NeuroImage*, *36*, 256–267.
- Han, S., Fan, Y., & Mao, L. (2008). Gender difference in empathy for pain: An electrophysiological investigation. *Brain Research*, *1192*, 85–93.
- Han, S., & Northoff, G. (2008). Culture-sensitive neural substrates of human cognition: A transcultural neuroimaging approach. *Nature Review Neuroscience*, *9*, 646–654.
- Han, S., Northoff, G., Vogeley, K., Wexler, B. E., Kitayama, S., & Varnum, M. E. W. (2013). A cultural neuroscience approach to the biosocial nature of the human brain. *Annual Review of Psychology*, *64*, 335–359.
- Harada, T., Li, Z., & Chiao, J. Y. (2010). Differential dorsal and ventral medial prefrontal representations of the implicit self-modulated by individualism and collectivism: An fMRI study. *Social Neuroscience*, *5*, 257–2571.
- Hein, G., Silani, G., Preuschoff, K., Batson, C. D., & Singer, T. (2010). Neural responses to ingroup and outgroup members' suffering predict individual differences in costly helping. *Neuron*, *68*, 149–160.
- Iacoboni, M. (2006). Failure to deactivate in autism: The co-constitution of self and other. *Trends in Cognitive Sciences*, *10*, 431–433.
- Jackson, P. L., Meltzoff, A. N., & Decety, J. (2005). How do we perceive the pain of others? A window into the neural processes involved in empathy. *NeuroImage*, *24*, 771–779.
- Konrath, S., Bushman, B. J., & Grove, T. (2009). Seeing my world in a million little pieces: Narcissism, self-construal, and cognitive-perceptual style. *Journal of Personality*, *77*, 1197–1228.
- Li, H. Z., Zhang, Z., Bhatt, G., & Yum, Y. (2006). Rethinking culture and self-construal: China as a middle land. *Journal of Social Psychology*, *146*, 591–610.
- Li, W., & Han, S. (2010). Perspective taking modulates event related potentials to perceived pain. *Neuroscience Letters*, *469*, 328–332.
- Lin, Z., & Han, S. (2009). Self-construal priming modulates the scope of visual attention. *Quarterly Journal of Experimental Psychology, Section A*, *62*, 802–813.
- Lin, Z., Lin, Y., & Han, S. (2008). Self-construal priming modulates visual activity underlying global/local perception. *Biological Psychology*, *77*, 93–97.
- Luo, S., Shi, Z., Yang, X., Wang, X., & Han, S. (in press). Reminders of mortality decrease midcingulate activity in response to others' suffering. *Social Cognitive and Affective Neuroscience*.
- Ma, Y., Bang, D., Wang, C., Allen, M., Frith, C., Roepstorff, A., & Han, S. (in press). Sociocultural patterning of neural activity during self-reflection. *Social Cognitive and Affective Neuroscience*.
- Ma-Kellams, C., & Blascovich, J. (2012). Inferring the emotions of friends versus strangers: The role of culture and self-construal. *Personality and Social Psychology Bulletin*, *38*, 933–945.
- Markus, H. R., & Kitayama, S. (1991). Culture and the self: Implication for cognition, emotion and motivation. *Psychological Review*, *98*, 224–253.

- Markus, H. R., & Kitayama, S. (2010). Cultures and selves: A cycle of mutual constitution. *Perspectives on Psychological Sciences*, 5, 420–430.
- Ma, Y., Wang, C., & Han, S. (2011). Neural responses to perceived pain in others predict real-life monetary donations in different socioeconomic contexts. *NeuroImage*, 57, 1273–1280.
- Meyer, M. L., Masten, C. L., Ma, Y., Wang, C., Shi, Z., Eisenberger, N. I., & Han, S. (2013). Empathy for the social suffering of friends and strangers recruits distinct patterns of brain activation. *Social Cognitive and Affective Neuroscience*, 8, 446–454.
- Obhi, S. S., Hogeveen, J., & Pascual-Leone, A. (2011). Resonating with others: The effects of self-construal type on motor cortical output. *Journal of Neuroscience*, 31, 14531–14535.
- Oyserman, D., Coon, H. M., & Kimmelmeier, M. (2002). Rethinking individualism and collectivism: Evaluation of theoretical assumptions and meta-analyses. *Psychological Bulletin*, 128, 3–72.
- Oyserman, D., & Lee, S. W. S. (2008). Does culture influence what and how we think? Effects of priming individualism and collectivism. *Psychological Bulletin*, 134, 311–342.
- Preston, S. D., & Hofelich, A. J. (2012). The many faces of empathy: Parsing empathic phenomena through a proximate, dynamic-systems view of representing the other in the self. *Emotion Review*, 4, 24–33.
- Sedikides, C., & Skowronski, J. J. (2009). Social cognition and self-cognition: Two sides of the same evolutionary coin? *European Journal of Social Psychology*, 39, 1245–1249.
- Sheng, F., & Han, S. (2012). Manipulations of cognitive strategies and intergroup relationships reduce the racial bias in empathic neural responses. *NeuroImage*, 61, 786–797.
- Sheng, F., Liu, Q., Li, H., Fang, F., & Han, S. (in press). Task modulations of racial bias in neural responses to others' suffering. *NeuroImage*.
- Sheng, F., Liu, Y., Zhou, B., Zhou, W., & Han, S. (2013). Oxytocin modulates the racial bias in neural responses to others' suffering. *Biological Psychology*, 92, 380–386.
- Singer, T., Seymour, B., O'Doherty, J., Stephan, K. E., Dolan, R. J., & Frith, C. D. (2006). Empathetic neural responses are modulated by the perceived fairness of others. *Nature*, 439, 466–469.
- Sui, J., & Han, S. (2007). Self-construal priming modulates neural substrates of self-awareness. *Psychological Science*, 18, 861–866.
- Sui, J., Hong, Y., Liu, C. H., Humphreys, G. W., & Han, S. (2013). Dynamic cultural modulation of neural responses to one's own and friend's faces. *Social Cognitive and Affective Neuroscience*, 8, 326–332.
- Sui, J., Liu, C. H., & Han, S. (2009). Cultural difference in neural mechanisms of self-recognition. *Social Neuroscience*, 4, 402–411.
- Thomsen, L., Sidanius, J., & Fiske, A. P. (2007). Interpersonal leveling, independence, and self-enhancement: A comparison between Denmark and the US, and a relational practice framework for cultural psychology. *European Journal of Social Psychology*, 37, 445–469.
- Varnum, M. E. W., Grossmann, I., Kitayama, S., & Nisbett, R. E. (2010). The origin of cultural differences in cognition: Evidence for the social orientation hypothesis. *Current Directions in Psychological Science*, 19, 9–13.
- Varnum, M. E. W., Shi, Z., Chen, A., Qiu, J., & Han, S. (in press). When "our" reward is the same as "my" reward: Self-construal priming shifts neural responses to own vs. friends' rewards. *NeuroImage*.
- Wang, C., Ma, Y., & Han, S. (in press). Self-construal priming modulates pain perception: Event-related potential evidence. *Cognitive Neuroscience*.
- Watson, P. J., Grisham, S. O., Trotter, M. V., & Bideman, M. D. (1984). Narcissism and empathy: Validity evidence for the narcissistic personality inventory. *Journal of Personality Assessment*, 48, 301–305.
- Xu, X., Zuo, X., Wang, X., & Han, S. (2009). Do you feel my pain? Racial group membership modulates empathic neural responses. *Journal of Neuroscience*, 29, 8525–8529.
- Zuo, X., & Han, S. (2013). Cultural experiences reduce racial bias in neural responses to others' suffering. *Culture and Brain*, 1, 34–46.